



PERFORMANCE, HAEMATOLOGY, AND SERUM BIOCHEMISTRY OF RABBIT BUCKS FED SUPPLEMENTAL LEVELS OF *SACCHAROMYCES CEREVISIAE*

I.P. Solomon¹, S. Mathew¹, E.F. Istifanus^{1*}, E.E. Tom¹

¹Department of Animal Science, Faculty of Agriculture, University of Uyo, Uyo, Akwa Ibom State, Nigeria

* Corresponding author e-mail: istifanusemanuel@yahoo.com

Abstract

This study was carried out to investigate the effect of *Saccharomyces cerevisiae* on the growth performance, haematological and serum biochemical indices of rabbit bucks. Eighteen (18) bucks were used for the study. They were assigned to three treatment groups with 6 bucks per treatment, replicated 3 times in a completely randomized design for twenty-eight (28) days. The bucks were allowed one week of acclimatization before the commencement of the study, during which they were fed commercial growers diets and forages. The bucks were fed a formulated ration supplemented with *S. cerevisiae* at 0.0, 0.3, and 0.6g, respectively, and were coded T1, T2, and T3, respectively. T1 served as control. Growth parameters, haematological indices, and serum chemistry were measured in the course of the study. The data collected were subjected to analysis of variance (ANOVA) using SPSS version 21, and means were separated using Duncan of the same software. The results revealed that *S. cerevisiae* did not have an effect ($p > 0.05$) on the growth parameters in the rabbit bucks. This study revealed that *Saccharomyces cerevisiae* did not have a significant effect ($p > 0.05$) on the packed cell volume (PCV) of bucks in the study, as T1, T2, and T3 all had similar ($p > 0.05$) PCV values of 39.00, 39.00, and 41.00% respectively. *S. cerevisiae* influenced ($p < 0.05$) serum enzymes but did not have a significant effect ($p > 0.05$) on total protein, blood urea, Alanine transaminase (ALT), creatinine, and glucose. In conclusion, dietary supplementation of *S. cerevisiae* at 0.60% did not have a significant effect on growth parameter or most haematological and serum biochemical indices.

Keywords: Yeast, rabbit bucks, *Saccharomyces cerevisiae*, probiotics, growth promoters

Introduction

The ban on the use of antibiotics as growth promoters in the EU has led to investigating different natural feed additives to replace dietary antibiotics (Mahrose et al. 2019). It is well known that high levels of antibiotics have been used in livestock and poultry production as growth promoters and for disease prophylaxis. However, a major problem with this practice is the occurrence of antibiotic residues in meat because of the wide use of antibiotics as well as antibiotic resistance in both humans and animals. It therefore, becomes imperative to search for new safe alternatives for improving growth performance, health and disease control in animals. Probiotics (bacterial and yeast cultures) are non-pathogenic microbial adjuncts, that have been used as feed supplements and also as growth promoters, improving the immune system of animals by promoting the composition and microbial balance in their guts (Besseboua and Ayad 2021). The probiotic characteristics of *Saccharomyces cerevisiae* have been documented

(Belhassen et al. 2016, Abd el-aziz et al. 2021, Besseboua and Ayad 2021, Adli et al. 2023). *Saccharomyces cerevisiae*, also known as baker's yeast, is rich in proteins, vitamins, and minerals (Piskur et al. 2006). The choice for *Saccharomyces cerevisiae* is based on availability in the study region, since all research is expected to address local problem(s).

However, there is a paucity of empirical studies on the effects of *S. cerevisiae* on rabbit bucks. Thus, this study was designed to evaluate the effect of *S. cerevisiae* on growth performance, haematological profile, and serum biochemical indices in rabbit bucks.

Materials and Methods

Experimental Site

The study was carried out at the Rabbitry Unit of the Teaching and Research Farm of the Department of Animal Science, University of Uyo, Uyo, Akwa Ibom State. Uyo is located at 5°2'N and 7°55'E, with a mean annual temperature of between 26 °C and 28 °C, while the mean annual rainfall ranges from 2000 mm to 3000 mm (Solomon and Udoh 2017).

Experimental Design

The study adopted a completely randomized design (CRD). Eighteen mongrel bucks, aged 20 - 24 weeks, were randomly assigned to three treatment groups. Each treatment was further replicated three times, with two bucks per replicate. The total period of the experiment was 28 days (4 weeks).

Animal and Management

The eighteen bucks were housed individually in a three-tier wooden hutch according to their treatments and managed under standard husbandry conditions with an ad libitum supply of feed and water. Other routine management practices were carried out in accordance with the best animal welfare principles.

Experimental Diet

The bucks were fed a formulated ration with the proportionate inclusion of *S. cerevisiae* in treatments two and three at 0.3 and 0.6 g, respectively, while treatment one, which served as the control, was not supplemented with *S. cerevisiae*. The diets were coded as T1, T2, and T3, respectively.

Data Collection

Growth Performance

To evaluate the growth performance of these bucks, their initial weights were recorded at the beginning of the experiment and thereafter on a weekly basis for the 4 weeks of the study. Other parameters evaluated, namely total and daily feed intake, total and daily weight gains, and feed conversion ratio (FCR).

Serum and Haematological parameters

Upon termination of the experiment, blood samples for biochemical and haematological analysis (2 mL each) were collected via the external ear veins of the bucks from each replicated. The samples for haematological analysis were collected into sample bottles containing ethylene diamine tetra acetic acid (EDTA) to prevent coagulation, while those for serum biochemistry were collected in plain sample bottles without EDTA. The blood samples were subjected to laboratory examinations using an auto analyser (Sysmex kx-21n). The serum blood samples were kept in a slope position until they were separated through centrifugation at 1000 rpm for 20 minutes, followed by cooling in a deep freezer. All samples were then taken to the laboratory within two hours after collection for analysis. The haematological parameters evaluated were packed cell volume (PCV), red blood cells (RBC), white blood cells (WBC), haemoglobin concentration (HGB), mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH), mean corpuscular haemoglobin concentration (MCHC), lymphocytes, eosinophils,

monocytes, neutrophils, and basophils, while serum biochemical indices measured were total protein, urea, creatinine, aspartate aminotransferase (AST), alkaline phosphatase (ALP), alanine amino transaminase (ALT), and glucose. Olorede et al. (1996) described the clinical routine procedures that were used to determine the serum biochemical indices.

Table 1. Composition of Experimental Grower Diet

Ingredient (%)	T1	T2	T3
	(0.00g <i>S. cerevisiae</i>)	(0.3g <i>S. cerevisiae</i>)	(0.6g <i>S. cerevisiae</i>)
Maize	35.50	35.50	35.50
SBM	19.00	19.00	19.00
Wheat offal	26.00	26.00	26.00
PKC	15.50	15.50	15.50
Bone meal	3.00	3.00	3.00
Premix	0.25	0.25	0.25
Common Salt	0.25	0.25	0.25
Lysine	0.25	0.25	0.25
Methionine	0.25	0.25	0.25
* <i>S. cerevisiae</i>	0.00	0.30	0.60
Total	100	100	100
Calculated Proximate Composition			
Crude Protein	17.11	17.11	17.11
Crude Fibre	6.65	6.65	6.65
Ether Extract	6.72	6.72	6.72
Calcium	1.15	1.15	1.15
Phosphorus	0.74	0.74	0.74
Lysine	1.00	1.00	1.00
ME	2644.20	2644.20	2644.20

Vitamin Premix supplied the following per kg of diet: vitamin A 10,000 i.u., vitamin D3 12,000 i.u., vitamin E 20,000 i.u., vitamin K 2.5 mg, thiamin 2.0mg, riboflavin 3.0mg, **S. cerevisiae*: *Saccharomyces cerevisiae*; SBM: Soybean Meal; PKC: Palm Kernel Cake; ME: Metabolizable Energy

Statistical Analysis:

The data collected were subjected to analysis of variance (ANOVA) using SPSS version 21, and means were separated using Duncan of the same software to separate significantly different means.

Results

Growth performance of rabbit bucks fed diets containing dietary levels of *Saccharomyces cerevisiae*

The results on the growth performance of rabbit bucks fed diets containing dietary levels of *S. cerevisiae* are presented in table 2. The results revealed that *S. cerevisiae* did not exhibit an effect ($p > 0.05$) on the growth parameters of the rabbit bucks. The final weight was similar ($p > 0.05$) in the study but was not significantly increased in T3. The addition of *S. cerevisiae* to the bucks' diets had no effect on total and daily weight gains, but it caused non-significant increases in groups treated with *S. cerevisiae* compared to the non-treated group. The values obtained in the current investigation study were 1772.67, 1778.67, and 1819 g for T1, T2, and T3, respectively, for final weights, and 389.33, 395.00 and 444.00 g for T1, T2, and T3, respectively, for the total

weight gain. Although there was no significant difference in the total and daily feed intake, rabbit bucks on T3 diets had an insignificantly ($p > 0.05$) lower feed intake. Bucks in T3 presented total feed intake of 2787.67 g, while T1 and T2 had total feed intake values of 3095.33 and 3204.00 g respectively. A non-significantly higher daily feed intake was observed in T2 ($p > 0.05$). Furthermore, better ($p > 0.05$) feed conversion in T3-treated bucks was noticed, but it remains insignificant ($p > 0.05$). FCR values in the study were 8.27 (T1), 8.12 (T2), and 6.41 (T3).

Table 2. Growth performance of rabbit bucks fed diets containing dietary levels of *Saccharomyces cerevisiae*

Parameters	T1	T2	T3	SEM
Initial weight (g)	1383.33	1383.67	1375.33	65.96
Final weight (g)	1772.67	1778.67	1819.33	71.64
Total weight gain (g)	389.33	395.00	444.00	32.31
Daily weight gain (g)	13.90	14.11	15.86	1.15
Feed intake (g)	3095.33	3204.00	2787.67	142.49
Daily feed intake (g)	110.55	114.43	99.56	5.09
Feed Conversion Ratio	8.27	8.12	6.41	0.46

SEM is the standard error of mean; means without letters were not significant ($p > 0.05$)

Haematological indices of rabbit bucks fed diets containing dietary levels of *Saccharomyces cerevisiae*

The dietary inclusion of *Saccharomyces cerevisiae* in the diets of rabbit bucks, as shown in Table 3, significantly affected white blood cells (WBC) ($p < 0.05$), platelets, and neutrophils in the bucks. However, other haematological parameters were not influenced by dietary levels of *Saccharomyces cerevisiae*. This study revealed that *Saccharomyces cerevisiae* did not have an impact in a significant manner on PCV ($p > 0.05$), as T1, T2, and, T3 all had similar ($p > 0.05$) PCV values of 39.00, 39.00, and, 41.00%, respectively.

A higher WBC was observed with inclusion of *Saccharomyces cerevisiae* in the diets of the bucks when compared with those without *Saccharomyces cerevisiae* in their diets (T1) ($p < 0.05$). Bucks fed T3 and T2 had WBC values 5.23 and $5.17 \times 10^9/\text{dL}$, respectively, while bucks on T1 diet had a WBC value of $3.80 \times 10^9/\text{dL}$. Significantly higher ($p < 0.05$) platelet was observed in bucks fed T2 in the study, while similar ($p > 0.05$) lower values were recorded in T1 and T3, respectively. *Saccharomyces cerevisiae* did not have an effect on RBC, as the values recorded were statistically similar ($p > 0.05$). The MCV, MCH, and MCHC were all similar ($p > 0.05$) across dietary treatment groups. Haemoglobin was not impacted by *Saccharomyces cerevisiae* in the study, as the values observed were 13.27, 13.10, and 13.38 g/dL, respectively, for T1, T2 and T3. Similarly, lymphocytes, eosinophils, basophils, and monocytes all showed no significant alteration ($p > 0.05$) with inclusion of *Saccharomyces cerevisiae* in the bucks' diets. However, neutrophils were substantially increased ($p < 0.05$) in bucks fed the T1 diet compared to those fed T2 and T3 diets respectively.

Table 3. Haematological indices of rabbit bucks fed diets containing dietary levels of *Saccharomyces cerevisiae*

Parameters	T1	T2	T3	SEM
Packed Cell Volume (%)	39.00	39.00	41.00	0.69
White Blood Cells ($\times 10^9/\text{dL}$)	3.80 ^b	5.17 ^a	5.23 ^a	0.29
Platelet	174.67 ^b	339.67 ^a	221.67 ^{ab}	30.61
Red Blood Cells ($\times 10^{12}/\text{L}$)	5.87	5.77	5.97	0.18
MCV (fl)	66.67	68.67	70.00	1.45

MCH (pg)	23.00	23.67	23.00	0.32
MCHC (%)	34.00	33.67	33.67	0.15
Haemoglobin (g/dL)	13.27	13.10	13.38	0.28
Lymphocytes (%)	47.00	59.00	59.00	2.71
Eosinophil (%)	2.67	2.67	2.00	0.53
Monocytes (%)	2.00	2.00	1.67	0.11
Neutrophils (%)	48.67 ^a	35.67 ^b	37.67 ^{ab}	2.67
Basophil (%)	0.00	1.00	0.00	0.17

MCV: mean corpuscular volume; MCH: Mean corpuscular haemoglobin; MCHC: Mean corpuscular haemoglobin concentration; SEM: Standard error of means; Means with different superscripts are significant ($p < 0.05$).

Serum biochemistry of rabbit bucks fed diets containing dietary levels of *Saccharomyces cerevisiae*

The result on serum biochemical indices of male rabbits fed different levels of dietary *S. cerevisiae* are presented in table 3. The present study showed that *S. cerevisiae* influenced serum enzymes ($P < 0.05$), but did not have any significant effect on total protein, blood urea, ALT, creatinine and glucose ($p > 0.05$). The values of total protein observed in the study were 60.67, 56.00, and 60.67 g/dL for T1, T2, and T3, respectively. Similar urea values were recorded for all treatments in the study ($p > 0.05$). Lower aspartate aminotransferase (AST) and alanine alkaline phosphatase (ALP) were detected when adding *S. cerevisiae* to the bucks diets ($p < 0.05$). AST was higher (168.67 μ L) in the control group (T1), while bucks on T2 diet presented the least AST value. Alkaline phosphatase was higher in bucks' diet that did not contain *S. cerevisiae*. Alanine aminotransaminase (ALT) did not show any significant statistical variation with dietary inclusion of *S. cerevisiae* in the bucks' diets ($p > 0.05$). Creatinine values were 154.00, 119.33, and 133.00 mg/dL for T1, T2, and T3, respectively. Glucose values were also similar across diets ($P > 0.05$). The values obtained in the study were 5.60, 5.55, and 6.30 g/dL for T1, T2, and T3, respectively.

Table 4. Serum biochemistry of rabbit bucks fed diets containing dietary levels of *Saccharomyces cerevisiae*

Parameters	T1	T2	T3	SEM
Total protein (g/dL)	60.67	56.00	60.67	1.03
Urea (mmol/L)	7.90	7.30	7.40	0.18
AST (μ L)	168.67 ^a	91.67 ^b	112.00 ^{ab}	15.00
ALP (μ L)	31.00 ^a	24.67 ^b	26.00 ^{ab}	1.13
ALT (μ L)	46.67	37.67	37.00	3.15
Creatinine (mg/dL)	154.00	119.33	133.00	8.17
Glucose (g/dL)	5.60	5.55	6.30	0.21

AST: Aspartate aminotransferase; ALP: Alkaline phosphatase; ALT: Alanine transaminase; SEM: Standard error of means; Means with different super Scripts are significant ($p < 0.05$).

Discussions

Growth performance of rabbit bucks fed diets containing dietary levels of *Saccharomyces cerevisiae*

The reported findings in this study showed that the diet with *S. cerevisiae* did not affect the weight of the animals. The reported findings are not in line with those of Besseboua and Ayad (2021), who noticed a decrease in the animals' weight. The insignificant effect of *S. cerevisiae* on the growth parameters in rabbit bucks confirmed earlier studies by Seyidoglu and Galip

(2014) and Belhassen et al. (2016), who reported no significant differences in the growth performance of rabbits fed diets supplemented with *S. cerevisiae*. Kimsé et al. (2012), in their investigation, noted that yeast did not affect final body weight, daily weight gain, or feed intake in New Zealand rabbits. Abdel-Aziz et al. (2021) also demonstrated insignificant improvement in the growth performance of mice administered *S. cerevisiae*. However, Saied et al. (2011), and Onwurah and Okejim (2014), reported significant improvements with *S. cerevisiae* supplementation in broiler chicks and broiler chickens, respectively. These variations may be attributed to the species of animals, age, dose, and environmental conditions.

Haematological indices of rabbit bucks fed diets containing dietary levels of *Saccharomyces cerevisiae*

The WBCs, platelets, and neutrophils of rabbit bucks fed diets supplemented with *S. cerevisiae* were significantly different in the study as compared to the control group, while other haematological parameters remained the same. This observation agrees with the reports of Seyidoglu and Galip (2014) and Belhassen et al. (2016), who noted that *S. cerevisiae* in the diets of rabbits had no effect on some haematological parameters, although they detected a slight increase in (PCV) and, haemoglobin (HGB) concentrations in rabbits supplemented with yeast, which differ with this study. The significant effect of *S. cerevisiae* supplementation in bucks' diets on WBC in the current study supports the reports of Paryad et al. (2008) and Mulatu et al. (2019), who recorded significantly elevated WBCs in their studies, which differs from the reported findings of Besseboua and Ayad (2021), for the established parameters. Mulatu et al. (2019) revealed that WBCs, packed cell volume (PCV) and haemoglobin (HGB) were higher in chickens fed diets containing *S. cerevisiae*. Elghandour et al. (2019) noticed that yeast-fed rabbits had more WBCs, and LCTs, compared to the rabbits fed the control diet. The significant increase in the WBC, platelets, and decrease in neutrophils suggests improved immunity, better clotting factors, and reduced parasitism in the bucks, supporting the earlier report of Kazuun and Kazuun (2019), who reported that probiotics have immune-stimulatory effects.

Serum biochemistry of rabbit bucks fed diets containing dietary levels of *Saccharomyces cerevisiae*

Total protein, urea, ALT, creatinine, and glucose were not influenced by *Saccharomyces cerevisiae* supplementation in the bucks' diets in the current study. Serum total protein is a marker of the synthetic function of the liver and a valuable guide to assess the severity of liver damage (Osigwe et al. (2017). The similarity, therefore, in the study suggests no liver damage in the bucks. Low or high total protein is an indication of liver disorders and malnutrition (Augustine et al. 2020). The insignificant effect of *S. cerevisiae* on ALT, urea, and creatinine in this study is similar to the findings of Abd El-aziz et al. (2021). Similar results were demonstrated by Seyidoglu and Galip (2014), who supplemented 3 g of yeast per kg in rabbit diet, and Ozsoy and Yalcin (2011) with *Saccharomyces cerevisiae* in broiler turkey. The reduction in AST and ALP may be triggered by the presence of *S. cerevisiae* in the bucks' diet. According to Alagbe and Adegbite (2019), serum enzyme values are triggered by the presence of anti-nutrients or toxic substances in the feed of animals. Raised ALT and AST have been identified by Yin and Tong (2014) to be biomarkers of hepatocellular damage, which is induced by these enzymes leaking into the blood stream. Hence, the results in this present investigation indicated that the hepatic functions of the bucks were not compromised in the course of the study.

Conclusions

The present findings showed that dietary supplementation of *S. cerevisiae* at 0.60% did not exhibit a significant effect on growth parameters or most haematological and serum biochemical indices in rabbit bucks, therefore implies that the inclusion of *S. cerevisiae* in the rabbit bucks diet may not be necessary since it has very minimal effect growth and most blood parameters add to the total cost of production.

References

- Abd el-aziz AH, Mahrose KHM, El-kasrawy NI, Alsenosy AA. 2021. Yeast as growth promoter in two breeds of growing rabbits with special reference to its economic implications. *Anais da academia brasileira de ciências*. 93(2):1-13.
- Ad DN, Sjöfjan O, Sholikin MM, Hidayat C, Utama DT, Jayanegara A, Puspita PS. 2023. The effects of lactic acid bacteria and yeast as probiotics on the performance, blood parameters, nutrient digestibility, and carcass quality of rabbits: A meta-analysis. *Italian journal of animal science*. 22(1):157-168.
- Alagbe JO, Adegbite MB. 2019. Haematological and serum biochemical indices of starter broiler chicks fed aqueous extract of *balanitesaegyptiaca* and *alchorneacordifolia* bark mixture. *International journal of biological, physical and chemical studies*. 1(1):8-15.
- Augustine C, Khobe D, Babakiri Y, Igwebuikue JU, Joel I, John T, Ibrahim A. 2020. Blood parameters of wistar albino rats fed processed tropical sickle pod (*senna obtusifolia*) leaf meal-based diets. *Translational Animal Science*. 4:778-782. doi: 10.1093/tas/txaa063.
- Barbara kazuñ, krzysztof kazuusing probiotics in freshwater larviculture 2019. *Fisheries and aquatic life*. 27:130-135.
- Belhassen T, Bonai A, Gerencsér Z, Matics Z, Tuboly T, Bergaoui R, Kovacs M. 2016. Effect of diet supplementation with live yeast *saccharomyces cerevisiae* on growth performance, caecal ecosystem and health of growing rabbits. *World Rabbit Science*. 24(3):191-200. doi: 10.4995/wrs.2016.3991
- Besseboua O, Ayad A. 2021. Effect of *saccharomyces cerevisiae* feed supplementation on haematology and reproductive parameters for Algerian rabbits. *Journal of applied life sciences and environment*. 2(186):111-122.
- Elghandour MMY, Tan ZL, Abu Hafsa SH, Adegbeyeye MJ, Greiner R, Ugbogu EA, Salem AZM. 2019. *Saccharomyces cerevisiae* as a probiotic feed additive to non and pseudo-ruminant feeding: a review. *Journal of Applied Microbiology*. 128(3):658-674. doi: 10.1111/jam.14416
- Kimsé M, Bayourthe C, Monteils V, Fortun-Lamothe L, Cauquil L, Combes S, Gidenne T. 2012. Live yeast stability in rabbit digestive tract: consequences on the caecal ecosystem, digestion, growth and digestive health. *Animal Feed science and Technology*. 173(3-4):235-243. Doi: 10.1016/j.anifeedsci.2012.01.012.
- Mahrose KHM, ALagawany M, Abd elhack ME, Mahgoub, SA, Attia FM. 2019. Influences of stocking density and dietary probiotic supplementation on growing japanese quail performance. *Anais da academia brasileira de ciências*. 91: e20180616. Doi 10.1590/0001-3765201920180479.
- Mulatu K, Ameha N, Girma M. 2019. Effects of feeding different levels of baker's yeast on performance and hematological parameters in broiler chickens. *Journal of World's Poultry Research*. 9(2):38-49. doi: 10.36380/jw pr.2019.5.
- Onwurah FB, Okejim JC. 2014. Effect of graded levels of baker's yeast (*saccharomyces cerevisiae*) in water on carcass and organ characteristics of broiler chickens. *Academic research international*. 5(5):128-133.
- Osigwe, CC, Akah PA, Nworu CS. 2017. Biochemical and haematological effects of the leaf extract of *Newbouldia laevis* in alloxan-induced diabetic rats. *Journal of biosciences and medicines*. 5:18-36.

- Özsoy B, Yalçın S. 2011. The effects of dietary supplementation of yeast culture on performance, blood parameters and immune system in broiler turkeys. *Ankara ünv. Vet. Fak. Derg.* 58:117-122. Doi: 10.1501/vetfak_0000002460.
- Paryard A, Mahmoudi M. 2008. Effect of different levels of supplemental yeast (*saccharomyces cerevisiae*) on performance, blood constituents and carcass characteristics of broiler chicks. *Journal of African Agricultural Research.* 3(12):835-842.
- Piškur J, Rozpędowska E, Polakova S, Merico A, Compagno C. 2006. How did *saccharomyces* evolve to become a good brewer? *Trends in genetics.* 22(4):183-186.
- Saied JM, Al-Jabary QH, Thalij KM. 2011. Effect of dietary supplement yeast culture on production performance and hematological parameters in broiler chicks. *International Journal of Poultry Science.* 10(5):376- 380. doi: 10.3923/ijps.2011.376.380.
- Seyidoğlu N, Galip N. 2014. Effects of *saccharomyces cerevisiae* and *spirulina platensis* on growth performances and biochemical parameters in rabbits. *Kafkas ünv. Vet. Fak. Derg.* 20(3):331-336. doi: 10.9775/kvfd.2013.9988.
- Solomon IP, Udoh UH. 2017. Haematological Indices in Three Genotypes (Naked Neck, Frizzled Feather, Normal Feathered) of Nigerian Local Chicken. *Advances in Life Science and Technology.* 54:29-34.
- SPSS. 2012. Statistical package for social science scientists. Version 21.0 for windows. IBM Corp., Armonk, NY.
- Yin IK, Tong KS. 2009. Elevated alt and ast in an asymptomatic person: what the primary care doctor should do? *Malaysian Family Physician: The Official Journal of the Academy of Family Physicians of Malaysia.* 4(2-3):98-99.